

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Prenatal Exposure to Perfluoroalkyl Substances and Adiposity in Early and Mid-Childhood

Ana María Mora, Emily Oken, Sheryl L. Rifas-Shiman, Thomas F. Webster, Matthew W.
Gillman, Antonia M. Calafat, Xiaoyun Ye, and Sharon K. Sagiv

Table of Contents

Supplemental Table/Figure

Table S1. Spearman correlation coefficients for prenatal plasma PFAS concentrations in children who had an early childhood assessment.

Table S2. Spearman correlation coefficients for adiposity measurements collected in mid-childhood.

Table S3. Associations of prenatal plasma PFAS concentrations with overweight and obesity in early and mid-childhood among all children and stratified by child sex.

Figure S1. Mean change (95% CI) in mid-childhood adiposity measurements associated with an IQR increment in prenatal plasma PFOA concentrations.

Table S4. Associations of prenatal plasma PFAS concentrations with overall and central adiposity measurements in early childhood among all children and stratified by child sex (models adjusted for all PFAS simultaneously).

Table S5. Associations of prenatal plasma PFAS concentrations with overall and central adiposity measurements in mid-childhood among all children and stratified by child sex (models adjusted for all PFAS simultaneously).

Table S6. Associations of prenatal plasma PFAS concentrations with overall and central

adiposity measurements in early childhood among all children and stratified by child sex, after adjusting for maternal GFR and albumin levels during pregnancy.

Table S7. Associations of prenatal plasma PFAS concentrations with overall and central adiposity measurements in mid-childhood among all children and stratified by child sex, after adjusting for maternal GFR and albumin levels during pregnancy.

Table S8. Longitudinal associations of prenatal plasma PFAS concentrations with overall and central adiposity measurements collected in early and mid-childhood, estimated using linear regression models with generalized estimation equations (GEE).
